Naegleria fowleri or Braineating amoeba

December 28, 2022

<u>In news</u>— South Korea has recently reported its first case of infection from *Naegleria fowleri* or "brain-eating amoeba".

About brain-eating amoeba-

- Naegleria is an amoeba, a single-celled organism, and only one of its species, called Naegleria fowleri, can infect humans.
- It was first discovered in Australia in 1965 and is commonly found in warm freshwater bodies, such as hot springs, rivers and lakes.
- The amoeba enters the human body through the nose and then travels up to the brain.
- This can usually happen when someone goes for a swim, or dive or even when they dip their head in a freshwater body.
- In some cases, it was found that people got infected when they cleaned their nostrils with contaminated water.
- Scientists haven't found any evidence of the spreading of Naegleria fowleri through water vapour or aerosol droplets.
- Once it goes to the brain, it destroys brain tissues and causes a dangerous infection known as primary amebic meningoencephalitis (PAM).
- So far, Naegleria fowleri has been found in all continents and declared as the cause of PAM in over 16 countries, including India.

What are the symptoms of PAM?

• The first signs of PAM start showing within one to 12 days after the infection.

- In the initial stages, they might be similar to symptoms of meningitis, which are headache, nausea and fever.
- In the later stages, one can suffer from a stiff neck, seizures, hallucinations, and even coma.
- The infection spreads rapidly and on average causes death within about five days.
- The fatality of PAM is as such that only four people have survived out of 154 known infected individuals in the United States from 1962 to 2021.

The treatment for the infection-

- As the Naegleria fowleri infection is rare and progresses quickly, scientists haven't been able to identify any effective treatments yet.
- At present, doctors treat it with a combination of drugs, including amphotericin B, azithromycin, fluconazole, rifampin, miltefosine, and dexamethasone.

Climate change impact on the spread of the infection-

- According to the US Centers for Disease Control and Prevention (CDC), with the rising global temperatures, the chances of getting Naegleria fowleri infection will go up as the amoeba mainly thrives in warm freshwater bodies.
- The organism best grows in high temperatures up to 46°C and sometimes can survive at even higher temperatures.
- Various recent studies have found that excess atmospheric carbon dioxide has led to an increase in the temperature of lakes and rivers.
- These conditions provide a more favourable environment for the amoeba to grow.
- Heat waves, when air and water temperatures may be higher than usual, may also allow the amoeba to thrive.